

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptacmb1647

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1			Web Page for STN Seminar Schedule - N. America
NEWS	2	NOV	21	CAS patent coverage to include exemplified prophetic substances identified in English-, French-, German-, and Japanese-language basic patents from 2004-present
NEWS	3	NOV	26	MARPAT enhanced with FSORT command
NEWS	4	NOV	26	CHEMSAFE now available on STN Easy
NEWS	5	NOV	26	Two new SET commands increase convenience of STN searching
NEWS	6	DEC	01	ChemPort single article sales feature unavailable
NEWS	7	DEC	12	GBFULL now offers single source for full-text coverage of complete UK patent families
NEWS	8	DEC	17	Fifty-one pharmaceutical ingredients added to PS
NEWS	9	JAN	06	The retention policy for unread STNmail messages will change in 2009 for STN-Columbus and STN-Tokyo
NEWS	10	JAN	07	WPIDS, WPINDEX, and WPIX enhanced Japanese Patent Classification Data
NEWS	11	FEB	02	Simultaneous left and right truncation (SLART) added for CERAB, COMPUAB, ELCOM, and SOLIDSTATE
NEWS	12	FEB	02	GENBANK enhanced with SET PLURALS and SET SPELLING
NEWS	13	FEB	06	Patent sequence location (PSL) data added to USGENE
NEWS	14	FEB	10	COMPENDEX reloaded and enhanced
NEWS	15	FEB	11	WTEXTILES reloaded and enhanced
NEWS	16	FEB	19	New patent-examiner citations in 300,000 CA/CAplus patent records provide insights into related prior art
NEWS	17	FEB	19	Increase the precision of your patent queries -- use terms from the IPC Thesaurus, Version 2009.01
NEWS	18	FEB	23	Several formats for image display and print options discontinued in USPATFULL and USPAT2
NEWS	19	FEB	23	MEDLINE now offers more precise author group fields and 2009 MeSH terms
NEWS	20	FEB	23	TOXCENTER updates mirror those of MEDLINE - more precise author group fields and 2009 MeSH terms
NEWS	21	FEB	23	Three million new patent records blast AEROSPACE into STN patent clusters
NEWS	22	FEB	25	USGENE enhanced with patent family and legal status display data from INPADOCDB
NEWS	EXPRESS	JUNE	27 08	CURRENT WINDOWS VERSION IS V8.3, AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.
NEWS	HOURS			STN Operating Hours Plus Help Desk Availability
NEWS	LOGIN			Welcome Banner and News Items
NEWS	IPC8			For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:40:16 ON 05 MAR 2009

=> file medline embase biosis caplus  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.22	0.22

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 12:40:41 ON 05 MAR 2009

FILE 'EMBASE' ENTERED AT 12:40:41 ON 05 MAR 2009  
Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:40:41 ON 05 MAR 2009  
Copyright (c) 2009 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 12:40:41 ON 05 MAR 2009  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (hydrophobic(w)interaction(w)chromatography or HIC) and (ammonium(w)acetate or CH3COONH4) and (ammonium(w)sulfate or ammonium(w)sulphate)

L1 20 (HYDROPHOBIC(W) INTERACTION(W) CHROMATOGRAPHY OR HIC) AND (AMMONIUM(W) ACETATE OR CH3COONH4) AND (AMMONIUM(W) SULFATE OR AMMONIUM(W) SULPHATE)

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 15 DUP REM L1 (5 DUPLICATES REMOVED)

=> s l2 and py<2004

L3 9 L2 AND PY<2004

=> dis ibib abs l3 1-9

L3 ANSWER 1 OF 9 MEDLINE on STN  
ACCESSION NUMBER: 2000259303 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 10797245  
TITLE: Purification of a cystic fibrosis plasmid vector for gene therapy using hydrophobic interaction chromatography.  
AUTHOR: Diogo M M; Queiroz J A; Monteiro G A; Martins S A; Ferreira G N; Prazeres D M  
CORPORATE SOURCE: Centro de Engenharia Biologica e Quimica, Instituto Superior Tecnico, Av. Rovisco Pais, 1000 Lisboa, Portugal.  
SOURCE: Biotechnology and bioengineering, (2000 Jun 5) Vol. 68, No. 5, pp. 576-83.  
Journal code: 7502021. ISSN: 0006-3592.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE) (RESEARCH SUPPORT, NON-U.S. GOV'T)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals

ENTRY MONTH: 200007  
ENTRY DATE: Entered STN: 14 Jul 2000  
Last Updated on STN: 10 Dec 2002  
Entered Medline: 6 Jul 2000

AB The success and validity of gene therapy and DNA vaccination in in vivo experiments and human clinical trials depend on the ability to produce large amounts of plasmid DNA according to defined specifications. A new method is described for the purification of a cystic fibrosis plasmid vector (pCF1-CFTR) of clinical grade, which includes an ammonium sulfate precipitation followed by hydrophobic interaction chromatography (HIC) using a Sepharose gel derivatized with 1,4-butanediol-diglycidylether. The use of HIC took advantage of the more hydrophobic character of single-stranded nucleic acid impurities as compared with double-stranded plasmid DNA. RNA, denatured genomic and plasmid DNAs, with large stretches of single strands, and lipopolysaccharides (LPS) that are more hydrophobic than supercoiled plasmid, were retained and separated from nonbinding plasmid DNA in a 14-cm HIC column. Anion-exchange HPLC analysis proved that >70% of the loaded plasmid was recovered after HIC. RNA and denatured plasmid in the final plasmid preparation were undetectable by agarose electrophoresis. Other impurities, such as host genomic DNA and LPS, were reduced to residual values with the HIC column (<6 ng/microg pDNA and 0.048 EU/microg pDNA, respectively). The total reduction in LPS load in the combined ammonium acetate precipitation and HIC was 400,000-fold. Host proteins were not detected in the final preparation by bicinchoninic acid (BCA) assay and sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) with silver staining. Plasmid identity was confirmed by restriction analysis and biological activity by transformation experiments. The process presented constitutes an advance over existing methodologies, is scaleable, and meets quality standards because it does not require the use of additives that usually pose a challenge to validation and raise regulatory concerns.  
Copyright 2000 John Wiley & Sons, Inc.

L3 ANSWER 2 OF 9 MEDLINE on STN  
ACCESSION NUMBER: 1986278562 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 3733935  
TITLE: Optimization of preparative hydrophobic interaction chromatographic purification methods.  
AUTHOR: Gooding D L; Schmuck M N; Nowlan M P; Gooding K M  
SOURCE: Journal of chromatography, (1986 May 30) Vol. 359, pp. 331-7.  
Journal code: 0427043. ISSN: 0021-9673.  
PUB. COUNTRY: Netherlands  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 198609  
ENTRY DATE: Entered STN: 21 Mar 1990  
Last Updated on STN: 21 Mar 1990  
Entered Medline: 16 Sep 1986

AB The chromatographic behavior of five proteins on hydrophobic interaction matrices having six different ligand arms was investigated using gradient elution with ammonium sulfate and ammonium acetate buffers at two pH values. The nature of the mobile phase and/or the ligand chain arm of the matrix was found to have substantial effect on the resolution, retention, and selectivity. Ovalbumin was moderately or highly retained with ammonium sulfate on all columns; however, with ammonium acetate, ovalbumin was not retained on SynChropak Hydroxypropyl and Propyl columns.

Chromatographic conditions developed for analytical hydrophobic interaction chromatography columns containing 6.5-micron packings were adapted to preparative columns packed with 30-micron SynChroprep packings for the separation of serum components. Dynamic load capacities were 4-13 mg of ovalbumin per ml of column volume.

L3 ANSWER 3 OF 9 EMBASE COPYRIGHT (c) 2009 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 1990283826 EMBASE  
TITLE: Evaluation of ammonium acetate as a volatile buffer for high-performance hydrophobic-interaction chromatography.  
AUTHOR: Konishi, T.; Kamada, M.; Nakamura, H.  
CORPORATE SOURCE: Kanto Chemical Co., Inc., 3-2-8 Nihonbashi Honcho, Chuo-ku, Tokyo 103, Japan.  
SOURCE: Journal of Chromatography, (1990) Vol. 515, pp. 279-283.  
ISSN: 0021-9673 CODEN: JOCRAM  
COUNTRY: Netherlands  
DOCUMENT TYPE: Journal; Conference Article; (Conference paper)  
FILE SEGMENT: 029 Clinical and Experimental Biochemistry  
LANGUAGE: English  
SUMMARY LANGUAGE: English  
ENTRY DATE: Entered STN: 13 Dec 1991  
Last Updated on STN: 13 Dec 1991

AB Hydrophobic-interaction chromatography (HIC) is a widely used technique for the separation of proteins without denaturation. In HIC, although, ammonium sulphate or sodium sulphate buffer is generally used as an eluent, volatile buffers such as ammonium acetate and ammonium formate seem to be advantageous in order to simplify the subsequent procedures including desalting. Therefore, the applicability of ammonium acetate buffer was evaluated, as a representative of volatile buffers for HIC, with respect to effects on the retention and peak broadening of proteins. Several proteins were successfully separated under the optimized conditions using volatile ammonium acetate buffer.

L3 ANSWER 4 OF 9 EMBASE COPYRIGHT (c) 2009 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 1987071421 EMBASE  
TITLE: Effects of mobile phase and ligand arm on protein retention in hydrophobic interaction chromatography.  
AUTHOR: Schmuck, M.N.; Nowlan, M.P.; Gooding, K.M.  
CORPORATE SOURCE: SynChrom, Inc., Lafayette, IN 47902, United States.  
SOURCE: Journal of Chromatography, (1986) Vol. Vol. 371, pp. 55-62.  
CODEN: JOCRAM  
COUNTRY: Netherlands  
DOCUMENT TYPE: Journal  
FILE SEGMENT: 029 Clinical and Experimental Biochemistry  
LANGUAGE: English  
ENTRY DATE: Entered STN: 11 Dec 1991  
Last Updated on STN: 11 Dec 1991

AB The retentive properties of a series of hydrophobic interaction chromatography packings with six different ligand arms (SynChropak Hydroxypropyl, Methyl, Propyl, Butyl, Pentyl, and Benzyl) were investigated with mobile phases of different ionic compositions and pH. Substitution of ammonium acetate for ammonium sulfate resulted in decreased retention for most combinations of proteins and ligands, although the retention of some proteins, such as lysozyme on the pentyl ligand, was unchanged by the salt substitution. Generally, lower pH resulted in reduced retention, but

the elution of lysozyme was more affected by pH than that of ovalbumin.

L3 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2003:76812 CAPLUS

DOCUMENT NUMBER: 138:131557

TITLE: Process involving cationic exchange chromatography and hydrophobic interaction chromatography for obtaining TGF $\beta$ , IGF-1, lactoperoxidase, and immunoglobulins from milk products

INVENTOR(S): Kivits, Marinus Gerardus Cornelis; Galama, Catharina Marina; Hendriks, Andor Wilhelm Joseph

PATENT ASSIGNEE(S): Campina B.V., Neth.; Numico Research B.V.

SOURCE: PCT Int. Appl., 26 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003008447	A1	20030130	WO 2002-NL496	20020722 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2454548	A1	20030130	CA 2002-2454548	20020722 <--
AU 2002318066	A1	20030303	AU 2002-318066	20020722 <--
AU 2002318066	B2	20071011		
EP 1409538	A1	20040421	EP 2002-747753	20020722
EP 1409538	B1	20090107		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
CN 1555384	A	20041215	CN 2002-818211	20020722
NZ 530704	A	20050729	NZ 2002-530704	20020722
AT 420108	T	20090115	AT 2002-747753	20020722
IN 2004CN00115	A	20051209	IN 2004-CN115	20040120
US 20040219225	A1	20041104	US 2004-484255	20040621
PRIORITY APPLN. INFO.:			EP 2001-202794	A 20010720
			EP 2001-202795	A 20010720
			WO 2002-NL496	W 20020722

AB The present invention relates to a process for extracting beneficial compds., in particular growth factors, such as TGF  $\beta$  and IGF-1 from milk. In this process a hydrophobic interaction chromatog. step is included. A resin having a Bu group, or a Ph group as the ligand is used as hydrophobic interaction resin. The resin can be eluted with a salt gradient which, when the ligand is a Ph group, contains substantially no alc., and thus resulting in fractions enriched in the desired growth factors. These fractions can be separated further by means of a hydroxyapatite column.

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:596149 CAPLUS

DOCUMENT NUMBER: 137:275156

TITLE: Influences of the mobile phase composition and temperature on the retention behavior of aromatic alcohol homologues in hydrophobic interaction chromatography

AUTHOR(S): Wei, Yinmao; Yao, Cong; Zhao, Jianguo; Geng, Xindu

CORPORATE SOURCE: Institute of Modern Separation Science, Northwest University, Xi'an, 710069, Peop. Rep. China

SOURCE: Chromatographia (2002), 55(11/12), 659-665  
CODEN: CHRGB7; ISSN: 0009-5893

PUBLISHER: Friedrich Vieweg & Sohn Verlagsgesellschaft mbH

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To eliminate the very complicated effects of chromatog. thermodyn. in hydrophobic interaction chromatog. (HIC) with biopolymers as solutes, homologs of neutral aromatic alcs. were selected as solutes for investigating their thermodyn. behavior in HIC. The effects of the mobile phase composition and temperature (0.apprx.80°) on the retention behavior of the homologs were studied extensively. The retention behavior of the homolog was characterized by the linear parameters in the stoichiometric displacement model for retention (SDM-R). The retention of small mols. is essentially controlled by non-specific interaction in HIC as well as in reversed phase liquid chromatog. (RPLC), and the parameters obtained were found to follow the homolog rule. Plots of the logarithm of retention of solutes in four kinds of salt solution vs. the reciprocal of the absolute temperature over a wide range were nonlinear, indicating a large heat capacity change associated with retention. The thermodyn. parameters demonstrate the retention of small mols. in HIC to be entropy-driven at low temperature and enthalpy-driven at high temperature

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 7 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2002:515174 CAPLUS

DOCUMENT NUMBER: 137:210089

TITLE: Studying the retention mechanism of hydrophobic interaction chromatography by using aromatic alcohol homologues as solute

AUTHOR(S): Wei, Yinmao; Zhao, Jianguo; Yao, Cong; Geng, Xindu

CORPORATE SOURCE: Institute of Modern Separation Science, Key Laboratory of Modern Separation Science in Shaanxi Province, Northwest University, Xi'an, 710069, Peop. Rep. China

SOURCE: Fenxi Huaxue (2002), 30(6), 641-644  
CODEN: FHHHDT; ISSN: 0253-3820

PUBLISHER: Zhongguo Huaxuehui "Fenxi Huaxue" Bianji Weiyuanhui

DOCUMENT TYPE: Journal

LANGUAGE: Chinese

AB The retention behaviors of aromatic alc. homologs in hydrophobic interaction chromatog. (HIC) were studied firstly. The retention of aromatic alc. conforms to homolog rule. However, the retention values increase first, and then decrease with the increase in the reciprocal of absolute temperature This relation between retention value and temperature can be expressed by the nonlinear Van't Hoff equation. The properties of aromatic alc. mols. were characterized by the linear parameters in stoichiometric displacement model for retention (SDM-R). The retention for small mols. in HIC is controlled in essential by the hydrophobic interaction force as well as in reversed phase liquid chromatog. (RPLC) and in HIC of biopolymer. Probably using small mols. as solute to study the retention mechanism of HIC is a new

reasonable way and probably lays a foundation to study the retention mechanism of small mols. and biopolymer in HIC.

L3 ANSWER 8 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:935765 CAPLUS  
DOCUMENT NUMBER: 136:50274  
TITLE: Method for isolating and purifying a protein based on microaggregation and adsorption on solid support and use of purified protein in therapeutics  
INVENTOR(S): Berna, Patrick; Clement, Christelle  
PATENT ASSIGNEE(S): Warner Lambert Company, USA; Meristem Therapeutics  
SOURCE: PCT Int. Appl., 46 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: French  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2001098473	A2	20011227	WO 2001-FR1985	20010622 <--
WO 2001098473	A3	20020502		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2810667	A1	20011228	FR 2000-8118	20000623 <--
FR 2810667	B1	20040903		
EP 1297116	A2	20030402	EP 2001-947593	20010622 <--
EP 1297116	B1	20060412		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2004505615	T	20040226	JP 2002-504622	20010622
AT 323154	T	20060415	AT 2001-947593	20010622
PRIORITY APPLN. INFO.:			FR 2000-8118	A 20000623
			WO 2001-FR1985	W 20010622

AB The invention concerns a method for isolating and purifying a protein of interest, in particular from a complex medium such as a plant extract Said method is characterized in that it comprises a step whereby a complex medium, comprising the solution containing the protein of interest to be purified and a solid support capable of enabling its adsorption, is brought in the presence of an agent capable of causing said protein to precipitate in soluble form.

The protein of interest is thus partly aggregated and adsorbed on the solid support without substantial formation of macro-aggregates in the solution capable of spontaneous elutriation. Thus, the method was applied to the isolation and purification of canine lipase from recombinant maize or tobacco. Ammonium sulfate was used to form microaggregates of the enzyme and the microaggregates were adsorbed to diatomaceous earth. The enzyme was further purified using ion-exchange and metal-chelate affinity chromatog.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 9 OF 9 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:781471 CAPLUS

DOCUMENT NUMBER: 135:328108  
 TITLE: Process and equipment for plasmid purification  
 INVENTOR(S): Nochumson, Samuel; Durland, Ross; Yu-speight, Audrey;  
 Welp, John; Wu, Kuoewi; Hayes, Rexford  
 PATENT ASSIGNEE(S): Valentis, Inc., USA  
 SOURCE: U.S. Pat. Appl. Publ., 15 pp., Cont. of U.S. Ser. No.  
 887,673, abandoned.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20010034435	A1	20011025	US 2001-774284	20010129 <--
US 7026468	B2	20060411		
US 20060106208	A1	20060518	US 2006-327987	20060109
PRIORITY APPLN. INFO.:			US 1996-22157P	P 19960719
			US 1997-887673	B1 19970703
			US 2001-774284	A1 20010129

AB A scalable alkaline lysis process, including procedures and devices for the isolation of large quantities (grams and kilograms) of plasmid DNA from recombinant E. coli cells is disclosed. Effective, controllable, and economical operation, and consistently low level of host chromosomal DNA in the final plasmid product result. The process involves a series of new unit operations and devices for cell resuspension, cell lysis, and neutralization. Thus, the RNA may be precipitated with high salt (1M KOAc and 7M NH4OAc) and the plasmid DNA may be purified by anion exchange chromatog. (with Fractogel EMD TMAE, for example) or by hydrophobic interaction chromatog. (e.g., with Octyl Sepharose 4 FF).

REFERENCE COUNT: 104 THERE ARE 104 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> FIL STNGUIDE  
 COST IN U.S. DOLLARS  
 FULL ESTIMATED COST

SINCE FILE ENTRY	TOTAL SESSION
59.42	59.64

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)  
 CA SUBSCRIBER PRICE

SINCE FILE ENTRY	TOTAL SESSION
-4.10	-4.10

FILE 'STNGUIDE' ENTERED AT 12:46:16 ON 05 MAR 2009  
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.  
 LAST RELOADED: Feb 27, 2009 (20090227/UP).

=> LOGOFF  
 ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF  
 LOGOFF? (Y)/N/HOLD:Y  
 COST IN U.S. DOLLARS

SINCE FILE ENTRY	TOTAL SESSION
0.21	59.85

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
------------	-------



CA SUBSCRIBER PRICE

ENTRY	SESSION
0.00	-4.10

STN INTERNATIONAL LOGOFF AT 12:47:55 ON 05 MAR 2009